

Mast Diagnostica GmbH Feldstrasse 20 DE-23858 Reinfeld Germany Tel: + 49 (0) 4533 2007 0 Fax: + 49 (0) 4533 2007 68 email: mast@mast-diagnostica.de Web: www.mast-group.com

# Tetrathionate Broth Base – DM219S

#### Introduction

MAST<sup>®</sup> Tetrathionate Broth Base is a selective liquid medium that can be used for the culture of *Salmonellae*. MAST<sup>®</sup> culture media is supplied in a dehydrated powder form, allowing the end-user to prepare a suitable medium for bacterial & fungal culture. It is suitable to be prepared in a variety of receptacles and at volumes that conform to the end-users desired purpose. The culture of bacterial and fungal species are essential for routine laboratory purposes.

FOR IN VITRO USE ONLY NOT FOR USE IN DIAGNOSIS OF HUMAN DISEASE

#### **Intended Purpose**

Tetrathionate Broth Base dehydrated culture medium powder is intended for use to produce a selective, fluid enrichment medium. When prepared in accordance with the instructions for use it produces a liquid medium for the selective enrichment of *Salmonellae*.

Tetrathionate Broth Base is intended to be used in conjunction with other *in vitro* tests. It is intended to be used by professional, trained, clinical laboratory users for *in vitro* use and is not intended for use in the diagnosis of disease or other conditions in humans or as the basis of treatment or case management decisions.

## Principle of the test

Culture media remains the gold standard for the growth and isolation of viable bacterial and fungal cells. The liquid culture medium is inoculated by sampling 3-5 independent colonies using a sterile loop and suspending in the prepared broth (broth suspension). Broth suspensions should be incubated under the appropriate atmospheric conditions and temperature for the target organisms(s), after which time the medium will become turbid as an indication of growth

These methods should be used in conjuction with other *in vitro* devices in the aid of diagnosis.

Once prepared a single aliquot of culture media broth is only for single use and cannot be re-used.

Mast Diagnostic

12 rue Jean-Jacques Mention CS91106, 80011 Amiens, CEDEX 1 France Tél: + 33 (0) 3 22 80 80 67 Fax: + 33 (0) 3 22 80 99 22 email: info@mast-diagnostic.fr Web: www.mast-group.com



#### Components

MAST<sup>®</sup> culture media is supplied in a dehydrated form for reconstitution by the end-user. The formulation of the product is described in Table 1.

#### Table 1. Formulation of DM219\*

Meterial Concentration in mediu	
Material	Concentration in medium
Peptone	5.0 g/L
Bile salts	1.0 g/L
Calcium carbonate	10.0 g/L
Sodium thiosulphate	30.0 g/L

\*Formulation may change to meet performance criteria.

The formulation is illustrative of the DM219 product range. The product is manufactured within an ISO:9001 and ISO:13485 environment. Inter-batch variation is expected to be minimal with no direct impact on the product.

#### Stability and storage

The expiry date applies to unopened containers of MAST<sup>®</sup> dehydrated culture media when stored in the primary container and in accordance with the manufacturer's instructions. The expiry date and batch number are indicated on each pack label.

- Store packs in a dry environment.
- Store packs at room temperature (10°C to 25°C).
- Avoid sources of moisture such as autoclaves, CO<sub>2</sub> incubators and water-baths.
- Limit the time a pack remains open whilst in use.
- This product is hygroscopic, avoid prolonged exposure to ambient moisture.
- For opened packs of dehydrated culture media ensure lid is firmly closed after every use.
- Before use ensure the appearance of the media conforms to the expected colour and texture i.e. free flowing, no excessive lumps. Media that is discoloured or lumpy should be further examined for performance against the recommended QC organism panel.

#### Warnings and precautions

- 1. Tetrathionate Broth Base is for *in vitro* use only, and must be used by trained professional laboratory staff.
- 2. All microbiological cultures and equipment used to transfer and manipulate them should be treated as infectious. Autoclave sterilise all biohazard waste before disposal in accordance with local regulations.
- 3. On receipt, store MAST<sup>®</sup> dehydrated culture media at the recommended storage temperature and conditions stated on the pack.
- 4. Do not store near sources of moisture or within high humidity environments.
- 5. Do not use if media powder is discoloured and/or lumpy, examine against recommended QC



Mast Diagnostica GmbH Feldstrasse 20 DE-23858 Reinfeld Germany Tel: + 49 (0) 4533 2007 0 Fax: + 49 (0) 4533 2007 68 email: mast@mast-diagnostica.de Web: www.mast-group.com

organism panel before continuing use. Discolouration could be a sign of degradation and must be examined further.

- 6. When handling the device ensure that local and regulatory health and safety advice is followed.
- 7. When handling the sterilised solution, beware of the temperature, use thermal resistant gloves where appropriate.
- 8. When preparing culture media after sterilisation, ensure that this is performed in an aseptic manner.

MAST<sup>®</sup> dehydrated culture media are supplied in a sealed primary container, which helps to prevent moisture ingress from the environment. The nature and frequency of use of the device is conducive to an end-user re-entering the container. When the product is not in use, the primary container should remain sealed.

## **Materials Provided**

Mast<sup>®</sup> dehydrated culture media is supplied in a powder form contained within a re-usable primary container for end-user reconstitution.

## Materials required but not provided

Standard microbiological supplies and equipment such as petri dishes, bottles, tubes, laminar flow cabinet, water bath, autoclave, balance, weigh boats, spatulas, thermometer, timer, additives such as defibrinated blood, deionised water, or suitable control strains of microorganisms.

## Procedure

- 1. Refer to pack label for quantities and volumes required. Prepare MAST<sup>®</sup> Tetrathionate Broth Base by suspending the powder in distilled or deionised water.
- Boil the suspension, cool to below 45°C and add 20ml of iodine solution.
- 3. Mix well and distribute 10ml amounts in test tubes.
- 4. After the addition of iodine solution the medium should be used on the same day. The base only will keep for several weeks at 2 to 8°C.

Refer to local Health and Safety handling procedures for infectious waste disposal guidelines.

## **Technical Guidance**

Observe the powder before use. If the powder is discoloured or lumpy, this could be a sign of degradation and must be further examined.

Mast Diagnostic 12 rue Jean-Jacques Mention CS91106, 80011 Amiens, CEDEX 1 France Tél: + 33 (0) 3 22 80 80 67 Fax: + 33 (0) 3 22 80 99 22 email: info@mast-diagnostic.fr Web: www.mast-group.com



## Interpretation of results

After incubation, the inoculated broth will become turbid as an indication of growth. The broth can then be plated onto the appropriate medium to confirm growth of target organisms.

## Limitations of use

MAST<sup>®</sup> media are not intended to be used as the sole, and primary isolation medium in instances where a failure to detect a pathogenic infection would result in death, serious illness or possible transmission of infectious disease.

# **Quality Control**

Check for signs of deterioration. Quality control must be performed with at least one organism to demonstrate expected performance. Do not use the product if the result with the control organism is incorrect. The list below illustrates a range of performance control strains which the end user can easily obtain.

#### Table 2. Suggested organisms for QC

Test Organisms	Result
<i>Salmonella typhimurium</i> ATCC <sup>®</sup> 14028	Growth
<i>Escherichia coli</i> ATCC <sup>®</sup> 25922	No growth

#### References

Bibliography is available on request.